

# Dynamic Prevalence of Sleep Disorder Following Stroke or transient Ischemic Attack

*By* Budi Susatia

ORIGINAL CONTRIBUTION

Dynamic Prevalence of Sleep Disorders Following Stroke or Transient Ischemic Attack

Systematic Review and Meta-Analysis

Faizul Hasan, MSN; Christopher Gordon, PhD; Dean Wu, PhD; Hui-Chuan Huang, PhD; Lia Taurussia Yuliana, BSc; Budi Susatia, MEd; Olyvia Freeska Dwi Marta, MSc; Hsiao-Yean Chiu, PhD

BACKGROUND AND PURPOSE: The exact prevalence of sleep disorders following stroke or transient ischemic attack (TIA) remains unclear. We aimed to determine the prevalence of sleep-disordered breathing, insomnia, periodic leg movement during sleep, and restless leg syndrome following stroke or TIA in acute, subacute, and chronic phases and examine the moderating effects of patient characteristics (eg, age) and methodological features (eg, study quality) on the prevalence.

METHODS: We performed a systematic review and meta-analysis. Embase and PubMed were searched from inception to December 18, 2019. We included 64 047 adults in 169 studies (prospective, retrospective, case-control, and cross-sectional study designs) reporting the prevalence of sleep disorders following stroke or TIA.

RESULTS: In the acute phase, the overall prevalence of mild, moderate, and severe sleep-disordered breathing was 66.8%, 50.3%, and 31.6% (95% CIs, 63.8–69.7, 41.9–58.7, and 24.9–39.1). In the subacute phase, the prevalence of mild, moderate, and severe sleep-disordered breathing was 65.5%, 44.3%, and 36.1% (95% CIs, 58.9–71.5, 36.1–52.8, and 22.2–52.8). In the chronic phase, the summary prevalence of mild, moderate, and severe sleep-disordered breathing was 66.2%, 33.1%, and 25.1% (95% CIs, 58.6–73.1, 24.8–42.6, and 10.9–47.6). The prevalence rates of insomnia in the acute, subacute, and chronic phases were 40.7%, 42.6%, and 35.9% (95% CIs, 31.8–50.3, 31.7–54.1, and 28.6–44.0). The pooled prevalence of periodic leg movement during sleep in the acute, subacute, and chronic phases was 32.0%, 27.3%, and 48.2% (95% CIs, 7.4–73.5, 11.6–51.7, and 33.1–63.5). The summary prevalence of restless leg syndrome in the acute and chronic phases was 10.4% and 13.7% (95% CIs, 6.4–16.4 and 2.3–51.8). Age, sex, comorbidities, smoking history, and study region had significant moderating effects on the prevalence of sleep disorders.

CONCLUSIONS: Sleep disorders following stroke or TIA are highly prevalent over time. Our findings indicate the importance of early screening and treating sleep disorders following stroke or TIA.

GRAPHIC ABSTRACT: An online graphic abstract is available for this article.

Key Words: meta-analysis ■ prevalence ■ restless leg syndrome ■ sleep ■ smoking

Globally, stroke is one of the leading causes of morbidity and mortality in adults. Stroke survivors often experience long-term physical and psychological problems that undermine their quality of life.

Sleep-disordered breathing (SDB) has been recognized as an independent risk factor for stroke. Notably, the American Heart Association and American Stroke

Association stroke guidelines have recommended considering sleep studies for the prevention of recurrent stroke or transient ischemic attack (TIA) on the basis of high prevalence in this population. Other sleep disorders, including insomnia, periodic leg movement during sleep (PLMS), and restless leg syndrome (RLS), have also been associated with an increased stroke risk.

Correspondence to: Hsiao-Yean Chiu, PhD, Taipei Medical University, 250 Wu-Hsing St, Taipei, 110 Taiwan. Email hychiu03152@tmu.edu.tw

The Data Supplement is available with this article at https://www.ahajournals.org/doi/suppl/10.1161/STROKEAHA.120.029847.

For Sources of Funding and Disclosures, see page XXX.

© 2021 American Heart Association, Inc.

Stroke is available at www.ahajournals.org/journal/str

## Nonstandard Abbreviations and Acronyms

<b>PLMS</b>	periodic leg movement during sleep
<b>RLS</b>	restless leg syndrome
<b>SDB</b>	sleep-disordered breathing
<b>TIA</b>	transient ischemic attack

Sleep disorders after stroke or TIA may impair optimal stroke rehabilitation and functional recovery.<sup>7,9</sup> Untreated poststroke sleep disorders can contribute to recurrent stroke.<sup>9</sup> However, the nature of sleep disorders following stroke has not been fully recognized. A wide variation in the prevalence may be one of the contributors to a lack of recognition in clinical and community-based settings. Hence, the clinical importance and effect of sleep disorders following stroke or TIA have recently gained attention.

Understanding the prevalence of sleep disorders following stroke or TIA can increase awareness and acknowledge its clinical relevance among health care providers and health policymakers. Consequently, health professionals could perform sleep screening earlier before or after stroke or TIA onset and designing effective treatments for relieving sleep disorders.

In this study, we comprehensively updated data on the prevalence of sleep disorders, including SDB, insomnia, PLMS, RLS, following stroke or TIA according to the different phases after stroke (acute, subacute, and chronic). Additionally, we examined the moderating effects of methodological features and patient characteristics on the prevalence of sleep disorders after stroke or TIA.

## METHODS

### Data Sources and Searches

The data of the current study are available from the corresponding author by reasonable request. This systematic review and meta-analysis were reported based on Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.<sup>10</sup> It was prospectively registered with PROSPERO (no. 55042019126422). PubMed and Embase were searched from their inception until December 18, 2019. The combinations of keywords used were as follows: ("stroke" OR "cerebrovascular accident" OR "transient ischemic attack" OR TIA) AND ("sleep" OR "sleep disorder" OR "circadian rhythm" OR "insomnia" OR "hypersomnia" OR "sleep apnea" OR "sleep apnoea" OR "sleep-wake cycle disorder" OR "narcolepsy" OR "sleep complaint" OR "sleep change" OR "sleep disturbance" OR "sleep disruption"). An example of the PubMed search strategy can be found in Table I in the Data Supplement. Additionally, the reference lists of all retrieved studies were manually searched for related studies.

84

### Study Selection

We included full-text studies that met the following criteria: (1) participants' mean age  $\geq 19$  years; (2) diagnosis of stroke or TIA; (3) reporting the outcome of the prevalence of SDB, insomnia, PLMS, or RLS following stroke or TIA for  $\geq 10$  participants; (4) prospective, cross-sectional, retrospective, or case-control study design; and (5) published or accepted for publication by a peer-reviewed journal. No language restriction was applied. Two investigators (F. Hasan and O.F.D. Marta) independently reviewed and screened titles and abstracts according to the eligibility criteria. Any disagreement was resolved through discussion with the third investigator (H.-Y. Chiu).

### Data Extraction and Assessment of Methodological Quality

Two reviewers (F. Hasan and O.F.D. Marta) separately extracted data, and disagreements were resolved through discussion. We classified the prevalence of sleep disorders following stroke and TIA as acute ( $<1$  month), subacute (1–3 months), and chronic ( $\geq 3$  months) phases.<sup>11</sup> Descriptions of outcome measures are listed in Table II in the Data Supplement.

The methodological quality of included studies was determined using the Critical Appraisal Checklist. Eight items were used to determine the quality of cross-sectional studies, 11 items were used for cohort studies, and 10 items for case-control studies.<sup>12</sup> Each item was rated as "yes," "no," "unclear," or "not applicable." Two researchers independently evaluated included studies (F. Hasan and O.F.D. Marta). Any disagreement was resolved by consensus.

### Data Synthesis and Analysis

All analyses were conducted using Comprehensive Meta-Analysis version 2.0 software (Biostat, Englewood, NJ). A random-effects model was used to estimate the prevalence with a 95% CI of sleep disorders following stroke or TIA in acute, subacute, and chronic phases. Between-study heterogeneity was assessed using Cochran  $Q$  and  $I^2$  tests. Substantial heterogeneity was determined if Cochran  $Q$  and  $I^2$  values were  $<0.05$  and at least 50%, respectively.<sup>13</sup> To identify possible reasons for heterogeneity, moderator, and metaregression analyses were performed. To ensure that sufficient data could be obtained for data analyses, moderator analyses were restricted to instances in which groups were represented by at least 2 studies. Egger intercept tests were used to test publication bias, and a  $P < 0.05$  represented a statistically significant publication bias.<sup>14</sup>


## RESULTS

### Selection, Inclusion, and Characteristics of Studies

We initially identified 7659 articles and screened abstracts and titles for inclusion (Figure 1). After removal of duplicates and irrelevant studies, 169 full-text articles were included in the review. Several studies reported more than one sleep outcome, resulting in that 132 included SDB, 28 insomnia, 10 PLMS, and 15 RLS. The characteristics of the included studies are listed in Tables

16

Downloaded from http://ahajournals.org by on January 7, 2021



**Figure 1. Study flow diagram.** n indicates number of studies; PLMS, periodic leg movement during sleep; RLS, restless leg syndrome; and SDB, sleep-disordered breathing.

through VI in the Data Supplement. Details about the inclusion and exclusion criteria of the included studies are listed in Table VII in the Data Supplement (including the reference list of the included studies).

For studies reporting SDB prevalence, the total sample size was 12 32 (stroke survivors, 12 971 and TIA cases, 1061); mean age was 64.9 years; and 63% were men. All studies were diagnosed as having SDB through in-laboratory (n=77) or home-based (n=55) polysomnography. Most of the studies considered obstructive and central sleep apnea as SDB and reported combined prevalence. Therefore, it is difficult to report the prevalence separately.

The studies examining insomnia had a total of 46 169 participants (stroke survivors, 46 133 and TIA cases,

36); the mean age was 63.0 years and 44.5% were women. Thirteen studies used the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, and the International Classification of Sleep Disorders-2 as references for the diagnosis of insomnia. Fifteen studies assessed insomnia symptoms by using self-administered questionnaires (7-item Insomnia Severity Index, 16-item Sleep Diagnosis Questionnaire, 32-item Sleep Habit Questionnaire, 7-item Hong Kong version of Insomnia Questionnaire, and 14-item St Mary's Sleep Questionnaire).

For PLMS prevalence, 10 studies involving 1165 stroke survivors and 32 TIA cases with a mean age of 63.2 years were included. All participants were diagnosed as having PLMS on the basis of

results of the periodic limb movement index derived from the in-laboratory (n=8) or home-based (n=2) polysomnography.

In terms of RLS, 15 studies involving 2639 patients (stroke survivors, 10 305 and TIA cases, 17) with a mean age of 63.4 years were included. Fourteen studies used the International Restless Legs Syndrome Study Group questionnaire for diagnosis of RLS, and one study used a single item to assess RLS.

## Prevalence of Sleep Disorders Following Stroke or TIA

### Sleep-Disordered Breathing

For the acute phase, the pooled prevalence of mild SDB was 66.8% (95% CI, 63.8–69.7), moderate SDB was 50.3% (95% CI, 41.9–58.7), and severe SDB was 31.6% (95% CI, 24.9–39.1; Figure 2A). For the subacute phase, 16, 13, and 9 included studies reported that the pooled prevalence of mild SDB was 65.5% (95% CI, 58.9–71.5), moderate SDB was 44.3% (95% CI, 36.1–52.8), and severe SDB was 36.1% (95% CI, 22.2–52.8; Figure 2A). For the chronic phase, the summary prevalence of mild SDB was 66.2% (95% CI, 58.6–73.1), moderate SDB was 33.1% (95% CI, 24.8–42.6), and severe SDB was 25.1% (95% CI, 10.9–47.6; Figure 2A).

### Insomnia

Overall, the pooled prevalence was 40.7% (95% CI, 31.8–50.3) for the acute phase, 42.6% (95% CI, 31.7–54.1) for the subacute phase, and 35.9% (95% CI, 28.6–44) for the chronic phase. Among these, studies reporting insomnia against standard references had a pooled prevalence of 32.5% (95% CI, 17.4–52.5) in the acute phase, 34.8% (95% CI, 20.6–52.4) in the subacute phase, and 37.1% (95% CI, 32.5–41.9) in the chronic phase. For self-reported insomnia symptoms, the summary prevalence was 47.1% (95% CI, 33.5–61.1) in the acute phase, 50.4% (95% CI, 38.4–62.3) in the subacute phase, and 36.9% (95% CI, 27.2–47.9) in the chronic phase (Figure 2B).

### Periodic Leg Movement Sleep Disorders

Ten studies reported the prevalence of PLMS. All studies included survivors of stroke with mild to moderate severity. The pooled prevalence in the acute phase was 32.0% (95% CI, 7.4–73.5), 27.3% (95% CI, 11.6–51.7) in the subacute phase, and 48.2% (95% CI, 33.1–63.5) in the chronic phase (Figure 2C).

### Restless Leg Syndrome

RLS prevalence was estimated in the acute phase after stroke in 13 studies and in the chronic phase in 2 studies. The summary prevalence of RLS was 10.4% in the acute phase (95% CI, 6.4–16.4) and 13.7% (95% CI, 2.3–51.8) in the chronic phase (Figure 2D).

## Heterogeneity Among Included Studies

As seen in Table VIII in the Data Supplement, considerable heterogeneity was identified among the included studies (all  $P < 0.001$ ).

## Subgroup and Metaregression Analyses

### Acute-Disordered Breathing

In the acute phase, age was positively associated with the pooled prevalence of mild SDB ( $B = 0.04$ ;  $P = 0.009$ ). Studies that included patients with comorbidities had a higher pooled prevalence of mild SDB than did those without comorbidities ( $P = 0.009$ , Table IX in the Data Supplement). Age, percentage of men in the study, body mass index, the country in which the study was conducted, comorbidities, and smoking were not significant moderators of prevalence in terms of moderate and severe SDB.

In the subacute phase, the studies that included patients with comorbidities yielded a higher prevalence of mild and severe SDBs than did those without comorbidities ( $P = 0.01$  and  $0.04$ , respectively).

In the chronic phase, the studies including patients with comorbidities yielded a higher prevalence of mild and moderate SDBs than did those without comorbidities ( $P = 0.02$  and  $0.002$ , respectively). Studies conducted in Eastern countries yielded a higher prevalence of severe SDB than did those conducted in Western countries ( $P = 0.009$ ).

### Insomnia


In the studies assessing insomnia, age, percentage of men in the study, the country in which the study was conducted, anxiety, depression, pain, comorbidities, and diagnosis were not significantly associated with insomnia prevalence (Table X in the Data Supplement).

### Periodic Leg Movement During Sleep

In the subacute phase, advancing age was associated with the pooled prevalence of PLMS ( $B = 0.48$ ;  $P < 0.001$ ; Table X in the Data Supplement). The studies, including patients with comorbidities, had a higher prevalence of PLMS than did those without comorbidities ( $P = 0.01$ ). Additionally, the studies including patients with the smoking habit exhibited a higher prevalence of PLMS than in patients without the smoking habit ( $P = 0.01$ ). In the chronic phase, only advancing age was positively associated with the pooled prevalence of PLMS ( $B = 0.07$ ;  $P < 0.001$ ).

### Restless Leg Syndrome

As shown in Table X in the Data Supplement, the percentage of men and body mass index correlated with the summary prevalence of RLS in the acute phase ( $B = -0.06$  and  $0.30$ , respectively;  $P = 0.02$  and  $0.01$ , respectively). Studies conducted in Western countries yielded a higher prevalence than did those conducted



**Figure 2.** Prevalence of sleep disorders with 95% CI, including (A) sleep-disordered breathing, (B) insomnia, (C) periodic leg movement sleep disorder, and (D) restless leg syndrome following stroke or transient ischemic attack during acute, subacute, and chronic phases. AHI indicates apnea-hypopnea index; and n, number of studies.

Downloaded from <http://ahajournals.org> by on January 7, 2021

in Eastern countries ( $P<0.001$ ). Because only 2 studies<sup>48</sup> in the chronic phase were included, we did not perform moderator and metaregression analyses.

### Publication Bias

The results<sup>60</sup> regard to publication bias are provided in Table XI in the Data Supplement. No publication bias was identified, except for mild SDB in the acute and chronic<sup>56</sup> phases ( $P=0.01$  and  $0.004$ , respectively<sup>56</sup>), insomnia in the chronic phase ( $P=0.03$ ), and RLS in the acute phase ( $P=0.01$ ).

### Assessment of Methodological Quality of Included Studies

The results of the Critical Appraisal Checklist of included studies are listed in Table XII in the Data Supplement.

## DISCUSSION

In the present meta-analysis, we examined the prevalence of SDB, insomnia, PLMS, and RLS in acute, subacute, and chronic phases following stroke or TIA. Previous meta-analyses have reported the prevalence of SDB and insomnia, but not PLMS and RLS, following stroke or TIA.<sup>15,16</sup> However, 2 meta-analyses of the prevalence of SDB did not further identify potential sources of high heterogeneity, thus limiting the internal validity of findings.<sup>15,16</sup> Furthermore, a meta-analysis<sup>17</sup> considered the prevalence of insomnia at different stroke phases as a whole because the number of studies included were few, which may have underestimated insomnia prevalence after stroke or TIA. Because we had a larger study sample, applied rigorous methodology, explored potential sources of heterogeneity, and analyzed prevalence rates according to different phases following stroke or TIA, our findings should be considered credible.<sup>110</sup>

The overall prevalence of mild, moderate, and severe SDB was gradually decreased in acute, subacute, and chronic phases, respectively. For example, in the acute phase, mild, moderate, and severe SDB prevalence rates gradually decreased from 66.8% to 50.3% to 31.6%, respectively. The downward trend was also observed in subacute (65.5%, 44.3%, and 36.1%, respectively) and chronic phases (66.2%, 33.1%, and 25.1%, respectively). In other words, the prevalence of mild SDB did not substantially change over time, whereas the prevalence of moderate and severe SDB appeared to decrease from acute to chronic phases, which is in line with previous reports.<sup>15,18,19</sup> The present study could not elucidate the cause of moderate and severe SDB prevalence reduction over time. Improvement in brain damage and chest function, decreased supine position duration, and recovery

from complications related to stroke may be attributed to the observed phenomenon.

Consistent with a previous publication,<sup>15</sup> our findings based on an additional 46 articles revealed that the prevalence of mild, moderate, and severe SDB was much higher in stroke survivors than in the general population (mild: 9%–38% and moderate to severe: 6%–17%).<sup>20</sup> However, the actual prevalence of SDB<sup>109</sup> was likely to be misestimated because some of the included studies did not exclude participants with sleep disorders before stroke. The associations between SDB and stroke are complex.<sup>21,22</sup> Accumulating evidence indicates that stroke can cause (or worsen preexisting) SDB through different mechanisms, such as damage to brain stem respiratory drive centers (eg, medullary lesion),<sup>23</sup> hypoglossal nerve dysfunction,<sup>24</sup> disturbed coordination of the upper airway, reduction of voluntary chest movement on the paralyzed side,<sup>22</sup> and prolonged supine position.<sup>25</sup> These may also play critical roles in the development of SDB after stroke or TIA.

The findings of our study revealed that age and comorbidities, which were also considered stroke contributors, were associated with the prevalence of SDB.<sup>26</sup> Decreased size of the upper airway lumen, pharyngeal airway lengthening, and change in the structures surrounding the pharynx<sup>27</sup> in older patients might be attributable to this phenomenon. Comorbidities related to sleep apnea (eg, chronic heart failure)<sup>30</sup> may also contribute to an increased prevalence of SDB, particularly central sleep apnea. We observed<sup>15</sup> that studies conducted in Eastern countries<sup>31–33</sup> had a higher prevalence of severe SDB in the chronic phase than those conducted in Western countries,<sup>34–36</sup> which presents a different perspective<sup>127</sup> on our existing knowledge. We did not find the role of body mass index and measure of obesity in the observation. Differences in the polysomnography setting may be a possible reason because we found that all included studies conducted in Eastern countries adopted in-laboratory polysomnography, whereas studies conducted in Western countries used home-based polysomnography. Moreover, attended portable monitoring is not recommended in the diagnosis of obstructive sleep apnea in adult patients with major comorbidities, owing to inadequate accuracy of such monitoring.<sup>37</sup>

In contrast to the previous meta-analysis<sup>17</sup> that estimated the overall prevalence of insomnia after stroke or TIA, our findings examined the pooled prevalence according to different phases after stroke and thus diagnostic methods. The prevalence of insomnia was found to be higher in patients with stroke or TIA than in the general population (6%–21%).<sup>38</sup> Although the cause of insomnia after stroke or TIA is not well understood, high stroke severity,<sup>39–41</sup> lesion location,<sup>42,43</sup> and negative emotions (ie, depression and anxiety) might play a role in the higher prevalence of insomnia following stroke or TIA.<sup>44</sup> In accordance with a prior work,<sup>45</sup> a slightly downward trend of overall insomnia prevalence was observed from

the subacute to chronic phases (42.6%–35.9%, respectively), supporting that insomnia is likely to subsequently recover over time following stroke or TIA. Neuronal regeneration and functional recovery after stroke may consolidate sleep continuity,<sup>46</sup> resulting in the improvement of insomnia.

Compared with the prevalence of PLMS in the general population (3.9%–28.6%),<sup>47–50</sup> stroke survivors were found to have a higher prevalence rate ranging from 27.3% to 48.2%. Possible mechanisms underlying the development of PLMS after stroke include white matter hyperintensities,<sup>51</sup> vascular brain damage,<sup>52</sup> and destructive brain lesions causing reticular formation on spinal pathways.<sup>53</sup> The pooled prevalence for PLMS after stroke or TIA exhibits an upward trend over time (32.0%–48.2%). Sleep consolidation may be attributed to the observation because we found that the prevalence of PLMS increased from the acute to the chronic phase, together with an expected improvement in insomnia. A consolidation in sleep continuity after the acute phase may result in more chances to produce a series of limb movements continuously.<sup>52</sup>

Age, presence of comorbidities, and smoking history were positively correlated with the prevalence of PLMS. PLMS can be found mostly in adults above 40 years of age.<sup>54,55</sup> Knowledge of potential mechanisms underlying this phenomenon is still limited; however, possible reciprocal influences between leg movements and the cyclic alternating pattern have been speculated.<sup>56,57</sup> In addition, evidence showed that Lewy body dementia was significantly correlated with PLMS in advancing age.<sup>58</sup> The association between comorbidities (eg, cardiovascular disease and renal disease) and elevated PLMS prevalence is not fully understood. Several reasons, such as sympathetic hyperactivity, peripheral vascular damage, and hypoxia, may partially explain the phenomenon in patients with cardiovascular diseases.<sup>59</sup> **107** Increased risk of iron deficiency anemia was reported as a risk factor for elevated PLMS.<sup>60</sup> Patients with chronic kidney disease can be at particularly high risk of RLS; additionally, the positive effect of smoking on PLMS prevalence is potentially mediated by stimulatory dopamine effects triggered by nicotine intake, which is similar to that suggested for Parkinson disease.<sup>61,62</sup>

Analysis of patients after stroke yielded a slightly higher prevalence estimation of RLS (acute and chronic phases, 10.4% and 13.7%, respectively) compared with that of **31** general population (3.9% and 14.3%, respectively, based on the International Restless Legs Syndrome Study Group criterion),<sup>50,63</sup> reflecting that **106** we may not be an aggravator of RLS. Coincidentally, a growing body of evidence has suggested RLS to be the first clinical manifestation related to cerebral infarction, such as brain stem and pontine infarction.<sup>64–66</sup> Additional studies are required to validate our findings.

Notably, women had a higher association with RLS prevalence after stroke. A growing body of evidence

72

has suggested that RLS may appear for the first time during pregnancy, and parity increases the risk of RLS later in life.<sup>67,68</sup> Our findings are in line with those of previous studies<sup>69,70</sup> wherein increased body mass index was identified **12** as a significant moderator of the prevalence of RLS. Possible explanations for the relationship between obesity and RLS include dysregulated dopamine metabolism, sympathetic hyperactivity, and **77** chronic inflammation in the brain.<sup>69,71</sup> Racial difference is an independent risk factor related **10** to the prevalence of RLS<sup>72,73</sup> because we found that studies conducted in Western countries yielded a higher prevalence than those conducted in Eastern countries. Our findings and currently available evidence suggest that RLS prevalence may vary depending on the specific population in question.

### Limitations and Strengths **92**

Several limitations of this meta-analysis must be acknowledged. First, some of the estimations of the prevalence of insomnia and RLS were obtained from a single question, which might result in **187** underreporting of intended sleep disorders. Second, because of the small number of included studies for the prevalence estimation of PLMS and RLS, our findings should be carefully interpreted. Additional investigations are warranted. Third, because only certain studies reported the exclusion of patients with sleep disorders before study enrollment, the effect of preexisting **105** disorders on our findings could not be avoided. Data on the prevalence of sleep disorders in African and Eastern populations, such as Russian and Indian populations, were limited. Fourth, we observed a statistical proof of publication bias in terms of mild SDB in acute and chronic phases, insomnia in **48** chronic phase, and RLS in acute phase. However, in this meta-analysis, we applied a rigorous study methodology and included a large sample size for credibility.

### Conclusions **54**

We suggest that sleep disorders are highly prevalent in stroke **73** survivors at different phases after stroke. Our findings indicate the clinical importance of early screening and treatment of sleep disorders following stroke or TIA. That is, sleep studies or sleep examinations should be performed earlier before or after stroke or TIA to provide primary or secondary preventive treatment. Additional prospective investigations that examine the prevalence and clinical effects of PLMS and RLS following stroke and TIA are required.

### ARTICLE INFORMATION **62**

Received March 18, 2020; final revision received November 8, 2020; accepted November 26, 2020.



**Affiliations**

School of Nursing, College of Nursing (F.H., H.-C.H., O.F.D.M., H.-Y.), Department of Neurology, School of Medicine, College of Medicine (D.W.), Department of Neurology, Shuang-Ho Hospital (D.W.), Research Center of Sleep Medicine, College of Medicine (D.W., H.-Y.C.), Taipei Medical University, Taiwan. School of Nursing (F.H., B.S.) and School of Midwifery (L.T.Y.), Politeknik Kesehatan Kemenkes Malang, Indonesia. Susan Wakil School of Nursing and Midwifery, Faculty of Medicine and Health, University of Sydney, Australia (C.G.). Nursing Department, Faculty of Health Science, University of Muhammadiyah Malang, Indonesia (O.F.D.M.).

**Sources of Funding**

This meta-analysis was supported by grants from the Ministry of Science and Technology, Taiwan (MOST 106-2314-B-038-058-MY3).

**Disclosures**

None.

**Supplemental Materials**

Tables I–XII  
References

**REFERENCES**

1. Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017;135:e146–e603. doi: 10.1161/CIR.12.000000000485
2. Chen Q, Cao C, Gong L, Zhang Y. Health related quality of life in stroke patients and risk factors associated with patients for return to work. *Medicine*. 2019;98.
3. Kim J, Kim Y, Yang KI, Kim DE, Kim SA. The relationship between sleep disturbance and functional status in mild stroke patients. *Ann Rehabil Med*. 2019;39:545–552. doi: 10.5535/arm.2015.39.4.545
4. Barone DA, Krieger AC. Stroke and obstructive sleep apnea: a review. *Curr Atheroscler Rep*. 2013;15:334. doi: 10.1007/s11883-013-0334-8
5. Kernan WN, Ovbiagele B, Black HR, Bravata DM, Chimowitz MI, Ezekowitz MD, Fang MC, Fisher M, Furie KL, Heck DV, et al; American Heart Association Stroke Council, Council on Cardiovascular and Stroke Nursing, Council on Clinical Cardiology, and Council on Peripheral Vascular Disease. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:2160–2366. doi: 10.1161/STR.000000000000024
6. Wu MP, Lin HJ, Weng SF, Ho CH, Wang JJ, Hsu YW. Insomnia subtypes and the subsequent risks of stroke: report from a nationally representative cohort. *Stroke*. 2014;45:1349–1354. doi: 10.1161/STROKEAHA.113.003675
7. Herrmann DM, Bassetti CL. Role of sleep-disordered breathing and sleep disturbances for stroke and stroke recovery. *Neurology*. 2017;87:1407–1416. doi: 10.1212/WNL.0000000000003037
8. Kop DL, Nam H, Thomas RJ, Yun CH. Sleep disturbances as a risk factor for stroke. *J Stroke*. 2018;20:12–32. doi: 10.5853/jos.201702887
9. Zhou M, Zhang X, Qu J. Coronavirus disease 2019 (COVID-19): a clinical update. *Front Med*. 2020;14:126–135. doi: 10.1007/s11684-020-0767-8
10. Moher D, Schulz KF, Simera I, Altman DG. Guidance for developers of health research reporting guidelines. *PLoS Med*. 2010;7:e1000217. doi: 10.1371/journal.pmed.1000217
11. Adams HP Jr, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE 3rd. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in acute stroke treatment. *Stroke*. 1993;24:35–41. doi: 10.1161/01.str.24.1.35
12. Moola S, Munn Z, Tufanaru C, Aromataris E, Sears K, Sfecu R, Currie M, Oureshi R, Mattis P, Lisy K, et al. Chapter 7: systematic reviews of etiology and risk. In: Aromataris E, Munn Z (eds). *JBIM Manual for Evidence Synthesis*. 2020. <https://synthesismanual.jbi.global>
13. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557–560. doi: 10.1136/bmj.327.7414.557
14. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315:629–634. doi: 10.1136/bmj.315.7109.629

15. Seiler A, Camilo M, Korostovtseva L, Haynes AG, Brill AK, Horvath T, Egger M, Bassetti CL. Prevalence of sleep-disordered breathing after stroke and TIA: meta-analysis. *Neurology*. 2019;10.1212/WNL.0000000000006904.
16. Johnson KG, Johnson DC. Frequency of sleep apnea in stroke and TIA patients: a meta-analysis. *J Clin Sleep Med*. 2010;6:131–137.
17. Baylan S, Griffiths S, Grant N, Broomfield NM, Evans JJ, Gardani M. Incidence and prevalence of post-stroke insomnia: a systematic review and meta-analysis. *Sleep Med Rev*. 2020;49:101222. doi: 10.1016/j.smrv.2019.101222
18. Bassetti CL, Milanova M, Gugger M. Sleep-disordered breathing and acute ischemic stroke: diagnosis, risk factors, treatment, evolution, and long-term clinical outcome. *Stroke*. 2006;37:967–972. doi: 10.1161/01.STR.0000208215.49243.c3
19. Parra O, Arboix A, Bechich S, García-Eroles L, Montserrat JM, López JA, Ballester E, Guerra JM, Sopena JJ. Time course of sleep-related breathing disorders in first-ever stroke or transient ischemic attack. *Am J Respir Crit Care Med*. 2000;161(2 Pt 1):375–380. doi: 10.1164/ajrcm.161.2.9903139
20. Senarathna CV, Perret JL, Lodge CJ, Lowe AJ, Campbell BE, Matheson MC, Hamilton GS, Dharmage SC. Prevalence of obstructive sleep apnea in the general population: a systematic review. *Sleep Med Rev*. 2017;34:70–81. doi: 10.1016/j.smrv.2016.07.002
21. Blissitt PA. Sleep-disordered breathing after stroke: nursing implications. *Stroke*. 2017;48:e81–e84. doi: 10.1161/STROKEAHA.116.013087
22. Alexiev F, Brill AK, Ott SR, Duss S, Schmidt M, Bassetti CL. Sleep-disordered breathing and stroke: chicken or egg? *J Thorac Dis*. 2018;10(suppl 4):S244–S252. doi: 10.21037/jtd.2018.12.66
23. Nogues M S, Heidele K, Benarroch E. Unilateral focal lesions in the rostralateral medulla influence chemosensitivity and breathing measured during wakefulness, sleep, and exercise. *J Neurol Neurosurg Psychiatry*. 2006;69:138–139. doi: 10.1136/jnnp.69.1.138
24. Brown DL, Chervin RD, Wolfe J, Hughes R, Concannon M, Lisabeth LD, Gruis KL. Hypoglossal nerve dysfunction and sleep-disordered breathing after stroke. *Neurology*. 2014;82:1149–1152. doi: 10.1212/WNL.000000000000263
25. Camilo MR, Fernandes RM, Sander HH, Nobre F, Santos-Pontelli T, Santos AC, Araujo DB, Leite JP, Pontes-Neto OM. Supine sleep and positional sleep apnea after acute ischemic stroke and intracerebral hemorrhage. *Clinics (Sao Paulo)*. 2012;67:1357–1360. doi: 10.61861/clinics/2012(12)02
26. Mérie C, Køber L, Olsen PS, Andersson C, Jensen JS, Torp-Pedersen C. Risk of stroke after coronary artery bypass grafting: effect of age and comorbidity. *Stroke*. 2012;43:38–43. doi: 10.1161/STROKEAHA.111.620880
27. Martin SE, Mathur R, Marshall I, Douglas NJ. The effect of age, sex, obesity and posture on upper airway size. *Eur Respir J*. 1997;10:2087–2090. doi: 10.1183/09031936.97.10092087
28. Shigeta Y, Enciso R, Ogawa T, Clark G. Changes in three dimensional simulation models of the airway which are due to increases in age or body mass index. *Respir Physiol Neurobiol*. 2008;162:5–6.
29. Malhotra A, Huang Y, Fogel R, Lasic S, Pillar G, Jakab M, Kikinis R, White DP. Aging influences on pharyngeal anatomy and physiology: the predisposition to pharyngeal collapse. *Am J Med*. 2006;119:72e9–72.14. doi: 10.1016/j.amjmed.2005.01.077
30. Vazir A, Hastings PC, Dayer M, McIntyre HF, Henein MY, Poole-Wilson PA, Cowie MR, Morrell MJ, Simonds AK. A high prevalence of sleep disordered breathing in men with mild symptomatic chronic heart failure due to left ventricular systolic dysfunction. *Eur J Heart Fail*. 2007;9:243–250. doi: 10.1111/jejheart.2006.08.001
31. Cho KH, Lee DH, Kim KM, Choi YH, Nam HS, Heo JH, Heo K, Kim YD. Relationship between sleep apnea and coronary artery calcium in patients with ischemic stroke. *Front Neurol*. 2019;10:819. doi: 10.3389/fnol.2019.00819
32. Chen CY, Hsu CC, Pei YC, Yu CC, Chen YS, Chen CL. Nocturia is an independent predictor of severe obstructive sleep apnea in patients with ischemic stroke. *J Neurol*. 2011;258:189–194. doi: 10.1007/s00415-010-5705-2
33. Srijithesh PR, Shukla G, Srivastava A, Goyal V, Singh S, Behari M. Validity of the Berlin Questionnaire in identifying obstructive sleep apnea syndrome when administered to the informants of stroke patients. *J Clin Neurosci*. 2018;340–343. doi: 10.1016/j.jocn.2010.04.048
34. Basile VS, Sundaram A, Hopyan JJ, et al. Identifying obstructive sleep apnea after stroke/TIA: evaluating four simple screening tools. *Sleep Med*. 2015;21:133–139. doi: 10.1016/j.sleep.2015.12.013
35. Cadilhac DA, Thorpe RD, Pearce DC, Barnes M, Rochford PD, Tarquinio N, Davis SM, Donnan GA, Pierce RJ; SCOPES II Study Group. Sleep disordered breathing in chronic stroke survivors. A study of the long term

follow-up of the SCOPES cohort: home based polysomnography. *J Neurosci*. 2005;12:632–637. doi: 10.1016/j.jocn.2004.08.014

36. Sacchetti ML, Di Mascio MT, Fiorelli M, Toni D, Roukos R, Minni A, Saponara M. Post stroke sleep apnea hypopnea syndrome: a series of 12 consecutive stroke cases. *Eur Rev Med Pharmacol Sci*. 2012;16:1295–1300.

37. Collop NA, Anderson WM, Boehlecke B, Claman D, Goldberg R, Gottlieb DJ, Hudgele D, Sateia M, Schwab R; Portable Monitoring Task Force of the American Academy of Sleep Medicine. Clinical guidelines for the use of unattended portable monitors in the diagnosis of obstructive sleep apnea in adult patients. Portable Monitoring Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2007;3:737–747.

38. Ohayon MM. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Med Rev*. 2002;6:97–111. doi: 10.1053/smv.2002.0186

39. Leppävuori A, Pohjasvaara T, Vataja R, Kaste M, Erkinjuntti T. Insomnia in ischemic stroke patients. *Cerebrovasc Dis*. 2002;14:90–97. doi: 10.1159/000064737

40. Joa KL, Kim WH, Choi HY, Park CH, Kim ES, Lee SJ, Kim SY, Ko SH, Jung HY. The effect of sleep disturbances on the functional recovery of rehabilitation inpatients following mild and moderate stroke. *Am J Phys Med Rehabil*. 2017;96:734–740. doi: 10.1097/PHM.0000000000000744

41. Palomäki H, Berg A, Meririnne E, Kaste M, Lönnqvist R, Lehtihalmes M, Lönnqvist J. Complaints of poststroke insomnia and its treatment with mianserin. *Cerebrovasc Dis*. 2003;15:56–62. doi: 10.1159/000067127

42. Chen X, Bi H, Zhang M, Liu H, Wang X, Zu R. Research of sleep disorders in patients with cerebral infarction. *J Stroke Cerebrovasc Dis*. 2015;24:2508–2513. doi: 10.1016/j.jstrokecerebrovasdis.2015.06.033

43. Koenigs M, Holliday J, Solomon J, Grafman J. Left dorsomedial frontal brain damage is associated with insomnia. *J Neurosci*. 2010;30:16041–16043. doi: 10.1523/JNEUROSCI.3745-10.2010

44. Roth T. Insomnia: definition, prevalence, etiology, and consequences. *J Clin Sleep Med*. 2007;3(5 suppl):S7–10.

45. Glozier N, Moullaali TJ, Sivertsen B, Kim D, Mead G, Jan S, Li Q, Hackett ML. The course and impact of poststroke insomnia in stroke survivors aged 18 to 65 years: results from the Psychosocial Outcomes In Stroke (POISE) Study. *Cerebrovasc Dis Extra*. 2017;7:9–20. doi: 10.1159/000455751

46. Kwakkel G, Buma FE, Selzer ME. Understanding the mechanisms underlying recovery after stroke. In: Kwakkel G, Cohen LG, Selzer ME, Miller RH, Clarke S, eds. *Textbook of Neural Repair and Rehabilitation: Volume 2: Medical Neurorehabilitation*. Vol 2. 2nd ed. Cambridge: Cambridge University Press; 2014:7–24.

47. Hornyak M, Feige B, Riemann D, Voderholzer U. Periodic leg movements in sleep and periodic limb movement disorder: prevalence, clinical significance and treatment. *Sleep Med Rev*. 2006;10:169–177. doi: 10.1016/j.smrv.2005.12.003

48. Scofield H, Roth T, Drake C. Periodic limb movements during sleep: population prevalence, clinical correlates, and racial differences. *Sleep*. 2008;31:1221–1227.

49. Haba-Rubio J, Martí-Soler H, Marques-Vidal P, Tobback N, Andries D, Preisig M, Waerber G, Vollenweider P, Kutalik Z, Tafti M, et al. Prevalence and determinants of periodic limb movements in the general population. *Ann Neurol*. 2016;79:464–474.

50. Ohayon MM, Roth T. Prevalence of restless legs syndrome and periodic limb movement disorder in the general population. *J Psychosom Res*. 2002;53:547–554. doi: 10.1016/s0022-3999(02)00443-9

51. Boulos MI, Murray BJ, Muir RT, Gao F, Szilagyi GM, Huroy M, Kiss A, Walters AS, Black SE, Lim AS, et al. Periodic limb movements and white matter hyperintensities in first-ever minor stroke or high-risk transient ischemic attack. *Sleep*. 2017;40:zsw080. doi: 10.1093/sleep/zsw080

52. Manconi M, Fanfulla F, Ferri R, Miano S, Haba-Rubio J, Heinzer R, Horvath T, Proserpio P, Young P, Moschovitis G, et al. Periodic limb movements during sleep in stroke/TIA: prevalence, course, and cardiovascular burden. *Neurology*. 2018;90:e1663–e1672. doi: 10.1212/WNL.00000000000005471

53. Benbir G, Karadeniz D. Periodic leg movements in sleep in patients with supratentorial cerebral infarction. *Acta Neurol Belg*. 2012;112:27–32. doi: 10.1007/s13760-011-0002-0

54. Sforza E, Juony C, Ibanez V. Time-dependent variation in cerebral and autonomic activity during periodic leg movements in sleep: implications for arousal mechanisms. *Clin Neurophysiol*. 2002;113:883–891. doi: 10.1016/s1388-2457(02)00066-4

55. Ferri R, Manconi M, Lanuzza B, Cosentino FI, Bruni O, Ferini-Strambi L, Zucconi M. Age-related changes in periodic leg movements during sleep in patients with restless legs syndrome. *Sleep Med*. 2008;9:790–798. doi: 10.1016/j.sleep.2007.08.020

56. Pennestri MH, Whittom S, Adam B, Petit D, Carrier J, Montplaisir J. PLMS and PLMW in healthy subjects as a function of age: prevalence and interval distribution. *Sleep*. 2006;29:1183–1187. doi: 10.1093/sleep/29.9.1183

57. Terzano MG, Parrino L, Spaggiari MC. The cyclic alternating pattern sequences in the dynamic organization of sleep. *Electroencephalogr Clin Neurophysiol*. 1988;69:437–447. doi: 10.1016/0013-4694(88)90066-1

58. Hibi S, Yamaguchi Y, Umeda-Kameyama Y, Yamamoto H, Iijima K, Momose T, Akishita M, Ouchi Y. The high frequency of periodic limb movements in patients with Lewy body dementia. *J Psychiatr Res*. 2012;46:1590–1594. doi: 10.1016/j.jpsychires.2012.07.007

59. Walters AS, Rye DB. Review of the relationship of restless legs syndrome and periodic limb movements in sleep to hypertension, heart disease, and chronic kidney disease. *World J Nephrol*. 2016;5:224–232. doi: 10.5277/wjn.v5.i3.224

60. Maung SC, El Sara A, Chapman C, Cohen D, Cukor D. Sleep disorders and chronic kidney disease. *World J Nephrol*. 2016;5:224–232. doi: 10.5277/wjn.v5.i3.224

61. Oksenberg A. Alleviation of severe restless legs syndrome (RLS) symptoms by cigarette smoking. *J Clin Sleep Med*. 2010;6:489–490.

62. Rye DB. Parkinson disease and RLS: the dopaminergic bridge. *Sleep Med*. 2004;5:317–328. doi: 10.1016/j.sleep.2004.01.016

63. Ohayon MM, O'Hara R, Vitiello MV. Epidemiology of restless legs syndrome: a synthesis of the literature. *Sleep Med Rev*. 2012;16:283–295. doi: 10.1016/j.smrv.2011.05.002

64. Ruppert E, Kilic-Huck U, Wolff V, Tatu L, Ghobadi M, Bataillard M, Bourgin P. Restless legs syndrome as a first manifestation of a cerebral infarct. *J Clin Sleep Med*. 2014;10:1037–1038. doi: 10.5664/jcsm.4028

65. Ruppert E, Kilic-Huck U, Wolff V, Tatu L, Lefebvre F, Chambe J, Rabin-Choteau L, Bataillard M, Bourgin P. Brainstem stroke-related restless legs syndrome: frequency and anatomical considerations. *Eur Neurol*. 2013;113–118. doi: 10.1159/000366416

66. Tuo HZ, Tian Cui YN, Ma XY, Xu CL, Bi HY, Zhang LY, Zhang YB, Le WD, Ondo W. Restless legs syndrome secondary to pontine infarction: clinical analysis of five cases. *Chronic Dis Transl Med*. 2017;3:186–190. doi: 10.1016/j.cjdtm.2017.08.001

67. Kim WH, Kim BS, Kim SK, Chang SM, Lee DW, Cho MJ, Bae JN. Restless legs syndrome in older people: a community-based study on its prevalence and association with major depressive disorder in older Korean adults. *Int J Geriatr Psychiatry*. 2012;27:565–572. doi: 10.1002/gps.2754

68. Innes KE, Selfe TK, Agarwal P. Prevalence of restless legs syndrome in North American and Western European populations: a systematic review. *Sleep Med*. 2011;12:623–634. doi: 10.1016/j.sleep.2010.12.018

69. Batool-Anwar S, Li Y, De Vito K, Malhotra A, Winkelman J, Gao X. Lifestyle factors and risk of restless legs syndrome: prospective cohort study. *J Clin Sleep Med*. 2016;12:187–194. doi: 10.5664/jcsm.5482

70. Gao X, Schwarzschild MA, Wang H, Ascherio A. Obesity and restless legs syndrome in men and women. *Neurology*. 2009;72:1255–1261. doi: 10.1212/WNL.0000345673.35676.1c

71. Allen R. Dopamine and iron in the pathophysiology of restless legs syndrome. *Sleep Med*. 2004;5:385–391. doi: 10.1016/j.sleep.2004.01.012

72. Alkhatzha A, Saeed A, Rashidzadeh W, Romaker AM. Racial differences in the prevalence of restless legs syndrome in a primary care setting. *Hosp Pract (1995)*. 2014;42:131–137. doi: 10.3810/hp.2014.08.1127

73. Sawanyawisuth K, Palinkas LA, Ancoli-Israel S, Dimsdale JE, Loreda JS. Ethnic differences in the prevalence and predictors of restless legs syndrome between Hispanics of Mexican descent and non-Hispanic Whites in San Diego County: a population-based study. *J Clin Sleep Med*. 2013;9:47–53. doi: 10.5664/jcsm.2338

# Dynamic Prevalence of Sleep Disorder Following Stroke or transient Ischemic Attack

ORIGINALITY REPORT

# 49%

SIMILARITY INDEX

## PRIMARY SOURCES

1	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet	139 words — 2%
2	<a href="http://tremorjournal.org">tremorjournal.org</a> Internet	134 words — 2%
3	<a href="http://academic.oup.com">academic.oup.com</a> Internet	129 words — 2%
4	<a href="http://stroke.ahajournals.org">stroke.ahajournals.org</a> Internet	112 words — 1%
5	<a href="http://www.karger.com">www.karger.com</a> Internet	110 words — 1%
6	<a href="http://boris.unibe.ch">boris.unibe.ch</a> Internet	88 words — 1%
7	Laetitia S. Gaspar, Cátia Sousa, Ana Rita Álvaro, Cláudia Cavadas, Alexandrina Ferreira Mendes. "Common risk factors and therapeutic targets in obstructive sleep apnea and osteoarthritis: an unexpectable link?", Pharmacological Research, 2020 Crossref	84 words — 1%
8	<a href="http://kwaas.org">kwaas.org</a> Internet	83 words — 1%
9	Carolina Lombardi, Gianfranco Parati, Davide Soranna, Antonella Zambon et al. "Periodic limb movements during sleep and blood pressure changes in sleep apnoea: Data	83 words — 1%

---

10	<a href="https://epdf.tips">epdf.tips</a> Internet	81 words — 1%
11	<a href="https://link.springer.com">link.springer.com</a> Internet	79 words — 1%
12	<a href="https://www.tandfonline.com">www.tandfonline.com</a> Internet	71 words — 1%
13	<a href="https://ir.ymlib.yonsei.ac.kr">ir.ymlib.yonsei.ac.kr</a> Internet	71 words — 1%
14	<a href="https://eurheartj.oxfordjournals.org">eurheartj.oxfordjournals.org</a> Internet	67 words — 1%
15	<a href="https://www.antisnoreshirt.com">www.antisnoreshirt.com</a> Internet	59 words — 1%
16	<a href="https://www.pmh.go.th">www.pmh.go.th</a> Internet	56 words — 1%
17	<a href="https://circheartfailure.ahajournals.org">circheartfailure.ahajournals.org</a> Internet	56 words — 1%
18	<a href="https://www.tobaccoinduceddiseases.org">www.tobaccoinduceddiseases.org</a> Internet	55 words — 1%
19	Chia-Rung Wu, Yu-Kang Tu, Li-Pang Chuang, Christopher Gordon et al. "Diagnostic meta-analysis of the Pediatric Sleep Questionnaire, OSA-18, and pulse oximetry in detecting pediatric obstructive sleep apnea syndrome", <i>Sleep Medicine Reviews</i> , 2020 Crossref	55 words — 1%
20	"Sleep Disorders Medicine", Springer Science and Business Media LLC, 2017 Crossref	55 words — 1%
21	<a href="https://www.researchgate.net">www.researchgate.net</a>	

---

Internet

54 words — 1%

22 [healthmetrics.heart.org](https://healthmetrics.heart.org)  
Internet

51 words — 1%

23 Krithika S. Rao, Jeffrey L. Spees. "Harnessing Epicardial Progenitor Cells and Their Derivatives for Rescue and Repair of Cardiac Tissue After Myocardial Infarction", *Current Molecular Biology Reports*, 2017  
Crossref

49 words — 1%

24 Patricia A. Blissitt. "Sleep-Disordered Breathing After Stroke", *Stroke*, 2017  
Crossref

46 words — 1%

25 [www.neurology.org](http://www.neurology.org)  
Internet

45 words — 1%

26 [www.dovepress.com](http://www.dovepress.com)  
Internet

45 words — 1%

27 [dmsjournal.biomedcentral.com](http://dmsjournal.biomedcentral.com)  
Internet

44 words — 1%

28 [www.med-sovet.pro](http://www.med-sovet.pro)  
Internet

43 words — 1%

29 Geert Mayer. "S2k guideline: insomnia in neurological disorders", *DGNeurologie*, 2020  
Crossref

42 words — 1%

30 [onlinelibrary.wiley.com](http://onlinelibrary.wiley.com)  
Internet

41 words — 1%

31 [escholarship.org](http://escholarship.org)  
Internet

39 words — 1%

32 [eu-ireland-custom-media-prod.s3-eu-west-1.amazonaws.com](http://eu-ireland-custom-media-prod.s3-eu-west-1.amazonaws.com)  
Internet

37 words — < 1%

33	<a href="http://www.europeanreview.org">www.europeanreview.org</a> Internet	36 words — < 1%
34	<a href="http://revistas.rcaap.pt">revistas.rcaap.pt</a> Internet	36 words — < 1%
35	<a href="http://neurovascular.fmrp.usp.br">neurovascular.fmrp.usp.br</a> Internet	36 words — < 1%
36	Shehzad Niazi, R. Robert Auger, Thomas D. Hurwitz. "An Overview of Restless Leg Syndrome for the Mental Health Professional", <i>Psychiatric Annals</i> , 2015 Crossref	35 words — < 1%
37	Shunjie Liu, Junhao Pan, Ke Tang, Qingfeng Lei, Lu He, Yangyang Meng, Xiaodong Cai, Zhong Li. "Sleep spindles, K-complexes, limb movements and sleep stage proportions may be biomarkers for amnesic mild cognitive impairment and Alzheimer's disease", <i>Sleep and Breathing</i> , 2019 Crossref	35 words — < 1%
38	<a href="http://jtd.amegroups.com">jtd.amegroups.com</a> Internet	35 words — < 1%
39	<a href="http://www.medrxiv.org">www.medrxiv.org</a> Internet	35 words — < 1%
40	Elisabeth Ruppert, Ulker Kilic-Huck, Valérie Wolff, Laurent Tatu et al. "Brainstem Stroke-Related Restless Legs Syndrome: Frequency and Anatomical Considerations", <i>European Neurology</i> , 2015 Crossref	33 words — < 1%
41	<a href="http://mafiadoc.com">mafiadoc.com</a> Internet	33 words — < 1%
42	<a href="http://www.scielo.br">www.scielo.br</a> Internet	32 words — < 1%
43	<a href="http://bmctoday.net">bmctoday.net</a> Internet	

32 words — < 1%

44 [mmj.com.ua](http://mmj.com.ua)  
Internet

31 words — < 1%

45 &NA; "Current World Literature :", Current Opinion  
in Neurology, 04/2001  
Crossref

30 words — < 1%

46 [clinicaltrials.gov](http://clinicaltrials.gov)  
Internet

30 words — < 1%

47 George D. Fulk, Pierce Boyne, Makenzie Hauger,  
Raktim Ghosh, Samantha Romano, Jonathan  
Thomas, Amy Slutzky, Karen Klingman. "The Impact of Sleep  
Disorders on Functional Recovery and Participation Following  
Stroke: A Systematic Review and Meta-Analysis",  
Neurorehabilitation and Neural Repair, 2020  
Crossref

29 words — < 1%

48 Hsiao-Yean Chiu, Pin-Yuan Chen, Li-Pang Chuang,  
Ning-Hung Chen, Yu-Kang Tu, Yu-Jung Hsieh, Yu-  
Chi Wang, Christian Guilleminault. "Diagnostic accuracy of the  
Berlin questionnaire, STOP-BANG, STOP, and Epworth  
sleepiness scale in detecting obstructive sleep apnea: A bivariate  
meta-analysis", Sleep Medicine Reviews, 2017  
Crossref

29 words — < 1%

49 Dayna A. Johnson, Sogol Javaheri, Na Guo, Cora  
L. Champion et al. "Objective Measures of Sleep  
Apnea and Actigraphy-Based Sleep Characteristics as Correlates  
of Subjective Sleep Quality in an Epidemiologic Study",  
Psychosomatic Medicine, 2020  
Crossref

28 words — < 1%

50 [www.ccjm.org](http://www.ccjm.org)  
Internet

28 words — < 1%

51 [trialsjournal.biomedcentral.com](http://trialsjournal.biomedcentral.com)  
Internet

28 words — < 1%

52 Alamin, Fathia Abd El Rahman Mohamed, and Caroline Edward Ayad. "Characterization of Upper Air Way Tract in Snoring and Non-Snoring Patients: A CT Based Study", Open Journal of Radiology, 2016. 27 words — < 1%

Crossref

53 Judy Triplett, David Ellis, Amber Braddock, Erin Roberts et al. "Temporal and region-specific effects of sleep fragmentation on gut microbiota and intestinal morphology in Sprague Dawley rats", Gut Microbes, 2020 27 words — < 1%

Crossref

54 [www.e-sciencecentral.org](http://www.e-sciencecentral.org) 27 words — < 1%

Internet

55 Hsiao-Yean Chiu, Hsin-Chien Lee, Pin-Yuan Chen, Ying-Fan Lai, Yu-Kang Tu. "Associations between sleep duration and suicidality in adolescents: A systematic review and dose-response meta-analysis", Sleep Medicine Reviews, 2018 27 words — < 1%

Crossref

56 A. Tefekli. "Peyronie's disease: a silent consequence of diabetes mellitus", Asian Journal of Andrology, 1/2006 25 words — < 1%

Crossref

57 [www.cureus.com](http://www.cureus.com) 24 words — < 1%

Internet

58 [pure.uva.nl](http://pure.uva.nl) 24 words — < 1%

Internet

59 [medicalonline.hu](http://medicalonline.hu) 24 words — < 1%

Internet

60 Giovanni Mariscalco, Stefano Rosato, Giuseppe F. Serraino, Daniele Maselli et al. "Prior Percutaneous Coronary Intervention and Mortality in Patients Undergoing Surgical Myocardial Revascularization", Circulation: Cardiovascular Interventions, 2018 22 words — < 1%

Crossref



61	<a href="http://www.koreascience.or.kr">www.koreascience.or.kr</a> Internet	21 words — < 1%
62	<a href="http://repub.eur.nl">repub.eur.nl</a> Internet	21 words — < 1%
63	<a href="http://neurotree.org">neurotree.org</a> Internet	20 words — < 1%
64	Vaishal S Shah, Vijay Sardana. "Study of Obstructive Sleep Apnea in Acute Ischemic Stroke Patients", Indian Journal of Sleep Medicine, 2020 Crossref	20 words — < 1%
65	<a href="http://www.eaeeg.org">www.eaeeg.org</a> Internet	20 words — < 1%
66	<a href="http://e-space.mmu.ac.uk">e-space.mmu.ac.uk</a> Internet	19 words — < 1%
67	<a href="http://archive.org">archive.org</a> Internet	18 words — < 1%
68	Ting-Jhen Chen, Yi-Wei Chung, Hui-Chen (Rita) Chang, Pin-Yuan Chen, Chia-Rung Wu, Shu-Hua Hsieh, Hsiao-Yean Chiu. "Diagnostic accuracy of the CAM-ICU and ICDSC in detecting intensive care unit delirium: A bivariate meta-analysis", International Journal of Nursing Studies, 2020 Crossref	18 words — < 1%
69	<a href="http://www.researchsquare.com">www.researchsquare.com</a> Internet	17 words — < 1%
70	Mollie McDermott, Devin L. Brown. "Sleep apnea and stroke", Current Opinion in Neurology, 2020 Crossref	17 words — < 1%
71	<a href="http://bmcnurs.biomedcentral.com">bmcnurs.biomedcentral.com</a> Internet	16 words — < 1%
72	<a href="http://worldwidescience.org">worldwidescience.org</a>	

Internet

16 words — < 1 %

73 [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet

16 words — < 1 %

74 Pamela S. Roberts, Shilpa Krishnan, Suzanne Perea Burns, Debra Ouellette, Monique R. Pappadis. "Inconsistent Classification of Mild Stroke and Implications on Health Services Delivery", Archives of Physical Medicine and Rehabilitation, 2020

Crossref

16 words — < 1 %

75 [www.nature.com](http://www.nature.com)

Internet

15 words — < 1 %

76 Samia Tasnim, Mariya Rahman, Priyanka Pawar, Xinli Chi et al. "Epidemiology of sleep disorders during COVID-19 pandemic: A systematic scoping review", Cold Spring Harbor Laboratory, 2020

Crossref Posted Content

14 words — < 1 %

77 "Sleep and Quality of Life in Clinical Medicine", Springer Science and Business Media LLC, 2008

Crossref

14 words — < 1 %

78 [www.thieme-connect.de](http://www.thieme-connect.de)

Internet

14 words — < 1 %

79 Hsiao-Yean Chiu, Hsin-Chien Lee, Pin-Yuan Chen, Ying-Fan Lai, Yu-Kang Tu. "Associations Between Sleep Duration and Suicidality in Adolescents: A Systematic Review and Dose-Response Meta-Analysis", Sleep Medicine Reviews, 2018

Crossref

13 words — < 1 %

80 [jamanetwork.com](http://jamanetwork.com)

Internet

13 words — < 1 %

81 [epdf.pub](http://epdf.pub)

Internet

12 words — < 1 %

82 [www.biomedcentral.com](http://www.biomedcentral.com)  
Internet

12 words — < 1 %

83 Amit Kumar, Rohit Kumar, Pranav Ish, Nitesh Gupta, Siddharth Raj Yadav, Shibdas Chakrabarti. "Sleep-disordered Breathing and Stroke: A Common yet Ignored Association", Indian Journal of Sleep Medicine, 2020  
Crossref

12 words — < 1 %

84 Ying Jiang, Qiwen Ben, Hong Shen, Weiqi Lu, Yong Zhang, Jun Zhu. "Diabetes mellitus and incidence and mortality of colorectal cancer: a systematic review and meta-analysis of cohort studies", European Journal of Epidemiology, 2011  
Crossref

11 words — < 1 %

85 [downloads.hindawi.com](http://downloads.hindawi.com)  
Internet

11 words — < 1 %

86 [bachlab.pitt.edu](http://bachlab.pitt.edu)  
Internet

11 words — < 1 %

87 Chiara Baglioni, Christoph Nissen, Adrian Schweinoch, Dieter Riemann et al. "Polysomnographic Characteristics of Sleep in Stroke: A Systematic Review and Meta-Analysis", PLOS ONE, 2016  
Crossref

10 words — < 1 %

88 D.-K. Kaynak. "Periodic leg movements in sleep and periodic limb movement disorder: a polysomnographic finding and/or a disorder?", Médecine du Sommeil, 2008  
Crossref

10 words — < 1 %

89 Xie, Wuxiang, Fanfan Zheng, and Xiaoyu Song. "Obstructive Sleep Apnea and Serious Adverse Outcomes in Patients with Cardiovascular or Cerebrovascular

10 words — < 1 %

Disease : A PRISMA-Compliant Systematic Review and Meta-Analysis", Medicine, 2014.

Crossref

90 Lei Guo, Yanan Liu, Lijun Liu, Shixiu Shao, Yanwei Cao, Jiaming Guo, Niu Haitao. "The CYP19A1 (TTTA)n repeat polymorphism, but not Arg264Cys polymorphism, may affect the risk of prostate cancer: evidence from a meta-analysis", Research Square, 2020  
10 words — < 1%  
Crossref Posted Content

91 orca.cf.ac.uk  
Internet  
9 words — < 1%

92 Zhen Fang. "Sleep-associated movement disorders and the risk of cardiovascular disease: A systematic review and meta-analysis", World Journal of Meta-Analysis, 2015  
9 words — < 1%  
Crossref

93 Minhui Liu, Susan M. McCurry, Basia Belza, Diana T. Buchanan, Adrian Dobra, Michael Von Korff, Michael V. Vitiello. "Effects of Pain, Insomnia, and Depression on Psychoactive Medication Supply in Older Adults With Osteoarthritis", Medical Care, 2018  
9 words — < 1%  
Crossref

94 Zhi-Xin Huang, Yan Huang, Jie Zeng, Hong Hao, Greg F. Petroski, Haike Lu, Xintong Liu, Zhenguo Liu. "Admission Glucose Levels May Increase the Risk for Early Neurological Deterioration in Females With Acute Ischemic Stroke", Frontiers in Neurology, 2020  
9 words — < 1%  
Crossref

95 www.aerzteblatt.de  
Internet  
9 words — < 1%

96 www.ajnr.org  
Internet  
9 words — < 1%

97 Susana Ferreira. "Prevalence and characteristics of sleep apnoea in patients with stable heart failure: Results from a heart failure clinic", BMC Pulmonary Medicine,  
9 words — < 1%

98	<a href="http://sub.edu.bd">sub.edu.bd</a> Internet	8 words — < 1%
99	<a href="http://repositorio.utad.pt">repositorio.utad.pt</a> Internet	8 words — < 1%
100	<a href="http://journals.sagepub.com">journals.sagepub.com</a> Internet	8 words — < 1%
101	<a href="http://academictree.org">academictree.org</a> Internet	8 words — < 1%
102	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet	8 words — < 1%
103	Alessandro Pincherle, Marta Pace, Simone Sarasso, Laura Facchin, Jens P. Dreier, Claudio L. Bassetti. "Sleep, Preconditioning and Stroke", Stroke, 2017 Crossref	8 words — < 1%
104	Kuan, Yi-Chun, Kuang-Wei Huang, Der-Jen Yen, Chaur-Jong Hu, Cheng-Li Lin, and Chia-Hung Kao. "Angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers reduced dementia risk in patients with diabetes mellitus and hypertension", International Journal of Cardiology, 2016. Crossref	8 words — < 1%
105	<a href="http://applicaties.csc.uu.nl">applicaties.csc.uu.nl</a> Internet	8 words — < 1%
106	"Disorders of Sleep and Circadian Rhythms in Parkinson's Disease", Springer Science and Business Media LLC, 2015 Crossref	8 words — < 1%
107	<a href="http://zombiedoc.com">zombiedoc.com</a> Internet	8 words — < 1%

- 
- 108 [www.sleepclinic.be](http://www.sleepclinic.be) 8 words — < 1%  
Internet
- 
- 109 Mark I Boulos, Trevor Jairam, Tetyana Kendzerska, James Im, Anastasia Mekhael, Brian J Murray. "Normal polysomnography parameters in healthy adults: a systematic review and meta-analysis", *The Lancet Respiratory Medicine*, 2019 8 words — < 1%  
Crossref
- 
- 110 J. A. Dempsey. "Pathophysiology of Sleep Apnea", *Physiological Reviews*, 01/01/2010 8 words — < 1%  
Crossref
- 
- 111 [www.essm.org](http://www.essm.org) 8 words — < 1%  
Internet
- 
- 112 [helda.helsinki.fi](http://helda.helsinki.fi) 8 words — < 1%  
Internet
- 
- 113 Claudio L. Bassetti. "Sleep and Stroke", Elsevier BV, 2017 8 words — < 1%  
Crossref
- 
- 114 [discovery.dundee.ac.uk](http://discovery.dundee.ac.uk) 8 words — < 1%  
Internet
- 
- 115 Carlos Berlanga-Macías, Diana P. Pozuelo-Carrascosa, Celia Álvarez-Bueno, Jose Alberto Martínez-Hortelano et al. "Relationship between exclusive breast feeding and cardiorespiratory fitness in children and adolescents: a protocol for a systematic review and meta-analysis", *BMJ Open*, 2018 8 words — < 1%  
Crossref
- 
- 116 Niko Männikkö, Heidi Ruotsalainen, Jouko Miettunen, Halley M Pontes, Maria Kääriäinen. "Problematic gaming behaviour and health-related outcomes: A systematic review and meta-analysis", *Journal of Health Psychology*, 2017 8 words — < 1%  
Crossref

117 [discovery.ucl.ac.uk](https://discovery.ucl.ac.uk)  
Internet

8 words — < 1%

118 Emelia J. Benjamin, Michael J. Blaha, Stephanie E. Chiuve, Mary Cushman, Sandeep R. Das, Rajat Deo, Sarah D. de Ferranti, James Floyd, Myriam Fornage, Cathleen Gillespie, Carmen R. Isasi, Monik C. Jiménez, Lori Chaffin Jordan, Suzanne E. Judd, Daniel Lackland, Judith H. Lichtman, Lynda Lisabeth, Simin Liu, Chris T. Longenecker, Rachel H. Mackey, Kunihiro Matsushita, Dariush Mozaffarian, Michael E. Mussolino, Khurram Nasir, Robert W. Neumar, Latha Palaniappan, Dilip K. Pandey, Ravi R. Thiagarajan, Mathew J. Reeves, Matthew Ritchey, Carlos J. Rodriguez, Gregory A. Roth, Wayne D. Rosamond, Comilla Sasson, Amytis Towfighi, Connie W. Tsao, Melanie B. Turner, Salim S. Virani, Jenifer H. Voeks, Joshua Z. Willey, John T. Wilkins, Jason HY. Wu, Heather M. Alger, Sally S. Wong, Paul Muntner. "Heart Disease and Stroke Statistics—2017 Update: A Report From the American Heart Association", *Circulation*, 2017  
Crossref

8 words — < 1%

119 [en.wikipedia.org](https://en.wikipedia.org)  
Internet

7 words — < 1%

120 Rong Ren, Guoping Huang, Jihui Zhang, Junying Zhou et al. "Age and severity matched comparison of gender differences in the prevalence of periodic limb movements during sleep in patients with obstructive sleep apnea", *Sleep and Breathing*, 2015  
Crossref

7 words — < 1%

121 Alastair J.S. Webb, Sara Mazzucco, Linxin Li, Peter M. Rothwell. "Prognostic Significance of Blood Pressure Variability on Beat-to-Beat Monitoring After Transient Ischemic Attack and Stroke", *Stroke*, 2018  
Crossref

7 words — < 1%

122 Siren E. Kouwenhoven, Caryl L. Gay, Linda N. Bakken, Anners Lerdal. "Depressive symptoms in

7 words — < 1%

acute stroke: A cross-sectional study of their association with sociodemographics and clinical factors", Neuropsychological Rehabilitation, 2013

Crossref

123 Naama Fund, Amit Green, Gabriel Chodick, Melanie Orin, Gideon Koren, Varda Shalev, Yaron Dagan. "The epidemiology of sleep disorders in Israel: results from a population-wide study", Sleep Medicine, 2020

Crossref

124 Scholle, Sabine, and Hans Christoph Scholle. "Leg movements and periodic leg movements during sleep in the development across childhood and adolescence from 1 to 18 years", Sleep Medicine, 2014.

Crossref

125 C. L. A. Bassetti, W. Randerath, L. Vignatelli, L. Ferini-Strambi et al. "EAN/ERS/ESO/ESRS statement on the impact of sleep disorders on risk and outcome of stroke", European Journal of Neurology, 2020

Crossref

126 Huang, Y.-H., N.-F. Chi, Y.-C. Kuan, L. Chan, C.-J. Hu, H.-Y. Chiou, and L.-N. Chien. "Efficacy of phenytoin, valproic acid, carbamazepine and new antiepileptic drugs on control of late-onset post-stroke epilepsy in Taiwan", European Journal of Neurology, 2015.

Crossref

127 Molly Mattsson, Deirdre Murray, Mairead Kiely, Fergus McCarthy, Elaine McCarthy, Regien Biesma, Fiona Boland. "Eating Behaviour, Physical Activity, TV Exposure and Sleeping Habits in Five Year Olds: A Latent Class Analysis", Research Square, 2020

Crossref Posted Content

128 [bmcpneurology.biomedcentral.com](https://www.bmcpneurology.biomedcentral.com)

Internet

129 Palimeris, Stephania. "Combining a Tailored Strength Training Program with Transcranial Direct-Current Stimulation (tDCs) to Improve Upper Extremity Function in Chronic



---

**130** Bruce Newman. "Iron depletion by whole-blood donation harms menstruating females: The current whole-blood-collection paradigm needs to be changed", *Transfusion*, 10/2006 6 words — < 1%  
Crossref

---

**131** [content.iospress.com:443](http://content.iospress.com:443) 6 words — < 1%  
Internet

---

EXCLUDE QUOTES      OFF  
EXCLUDE BIBLIOGRAPHY      OFF

EXCLUDE MATCHES      OFF